

CONFORMAL HIGGS, OR TECHNI-DILATON - COMPOSITE HIGGS NEAR CONFORMALITY

KOICHI YAMAWAKI

*Department of Physics, Nagoya University
Nagoya, Japan**

In contrast to the folklore that Technicolor (TC) is a “Higgsless theory”, we shall discuss existence of a composite Higgs boson, Techni-Dilaton (TD), a pseudo-Nambu-Goldstone boson of the scale invariance in the Scale-invariant/Walking/Conformal TC (SWC TC) which generates a large anomalous dimension $\gamma_m \simeq 1$ in a wide region from the dynamical mass $m = \mathcal{O}(\text{TeV})$ of the techni-fermion all the way up to the intrinsic scale Λ_{TC} of the SWC TC (analogue of Λ_{QCD}), where Λ_{TC} is taken typically as the scale of the Extended TC scale Λ_{ETC} : $\Lambda_{\text{TC}} \simeq \Lambda_{\text{ETC}} \sim 10^3 \text{ TeV} (\gg m)$. All the techni-hadrons have mass on the same order $\mathcal{O}(m)$, which in SWC TC is extremely smaller than the intrinsic scale $\Lambda_{\text{TC}} \simeq \Lambda_{\text{ETC}}$, in sharp contrast to QCD where both are of the same order. The mass of TD arises from the *non-perturbative scale anomaly* associated with the techni-fermion mass generation and is typically 500-600 GeV, even *smaller than other techni-hadrons* of the same order of $\mathcal{O}(m)$, in another contrast to QCD which is believed to have no scalar $\bar{q}q$ bound state lighter than other hadrons. We discuss the TD mass in various methods, Gauged NJL model via ladder Schwinger-Dyson (SD) equation, straightforward calculations in the ladder SD/ Bethe-Salpeter equation, and the holographic approach including techni-gluon condensate. The TD may be discovered in LHC.

Keywords: Walking Technicolor, Scale Invariance, Conformal Symmetry, Techni-Dilaton, Fixed Point, Composite Higgs, Large Anomalous Dimension, Holographic Gauge Theory

1. Introduction

Toshihide Maskawa is famous for 2008 Nobel prize-winning paper with Makoto Kobayashi on CP violation but did also fundamental contributions particularly to the SCGT, the topics of this workshop: Back in 1974 he found with Hideo Nakajima¹ that spontaneous chiral symmetry breaking ($S\chi SB$) solution does exist for and only for the strong gauge coupling, with the critical coupling of order 1, based on the ladder Schwinger-Dyson (SD) equation with non-running (scale-invariant) coupling, namely the walking gauge dynamics what is called today. This turned out to be the origin of SCGT activities toward understanding the Origin of Mass. The present workshop SCGT 09 was held in honor of his 70th birthday on February 7, 2010 and the 35th anniversary of his crucial contributions to SCGT. I will later explain

*Present Address: Kobayashi-Maskawa Institute for the Origin of Particles and the Universe (KMI), Nagoya University. E-mail: yamawaki@kmi.nagoya-u.ac.jp

impact of Maskawa-Nakajima solution on the conformal gauge dynamics.

The Origin of Mass is the most urgent issue of the particle physics today and is to be resolved at the LHC experiments. In the standard model (SM), all masses are attributed to a single parameter of the vacuum expectation value (VEV), $\langle H \rangle$ of the hypothetical elementary particle, the Higgs boson. The VEV simply picks up the mass scale of the input parameter M_0 which is tuned to be tachyonic ($M_0^2 < 0$) in such a way as to tune $\langle H \rangle \simeq 246 \text{ GeV}$ (“naturalness problem”). As such SM does not explain the Origin of Mass.

Technicolor (TC)² is an attractive idea to account for the Origin of Mass without introducing ad hoc Higgs boson and tachyonic mass parameter: The mass arises *dynamically* from the condensate of the techni-fermion and the anti techni-fermion pair $\langle \bar{T}T \rangle$ which is triggered by the attractive gauge forces between the pair analogously to the quark-antiquark condensate $\langle \bar{q}q \rangle$ in QCD. For the TC with $SU(N_{\text{TC}})$ gauge symmetry and N_f flavors ($N_f/2$ weak doublets) of techni-fermions, the techni-pion decay constant $F_\pi = \langle H \rangle / \sqrt{N_f/2}$ corresponds to the pion decay constant $f_\pi \simeq 93 \text{ MeV}$ in QCD, and hence the TC may be a scale-up of QCD by the factor $F_\pi/f_\pi \simeq 2650/\sqrt{N_f/2}$. Then the mass scale of the condensate $\Lambda_\chi = (-\langle \bar{T}T \rangle / N_{\text{TC}})^{1/3}$ as the Origin of Mass may be estimated as

$$\Lambda_\chi \simeq \left(\frac{-\langle \bar{q}q \rangle}{N_c} \right)^{1/3} \cdot \frac{F_\pi / \sqrt{N_{\text{TC}}}}{f_\pi / \sqrt{N_c}} \simeq 450 \text{ GeV} \cdot \left(\frac{N_c / N_{\text{TC}}}{N_f / 2} \right)^{1/2}, \quad (1)$$

where we have used a typical value $(-\langle \bar{q}q \rangle)^{1/3} \simeq 250 \text{ MeV}$ ($N_c = 3$).

The dynamically generated mass scale of the condensate Λ_χ , or the dynamical mass of the techni-fermion, m ($\sim \Lambda_\chi \sim F_\pi$), in fact picks up the intrinsic mass scale Λ_{TC} of the theory (analogue of Λ_{QCD} in QCD) already generated by the scale anomaly through quantum effects (“dimensional transmutation”) in the gauge theory which is scale-invariant at classical level (for massless flavors):

$$\Lambda_{\text{TC}} = \mu \cdot \exp \left(- \int^{\alpha(\mu)} \frac{d\alpha}{\beta(\alpha)} \right) = \Lambda_0 \cdot \exp \left(- \int^{\alpha(\Lambda_0)} \frac{d\alpha}{\beta(\alpha)} \right), \quad (2)$$

where the running of the coupling constant $\alpha(\mu)$, with non-vanishing beta function $\beta(\alpha) \equiv \mu \frac{d\alpha(\mu)}{d\mu} \neq 0$, is a manifestation of the scale anomaly and Λ_0 is a fundamental scale like Planck scale. Note that Λ_{TC} is independent of the renormalization point μ , $\frac{d\Lambda_{\text{TC}}}{d\mu} = 0$, and can largely be separated from Λ_0 through logarithmic running (“naturalness”). Thus the Origin of Mass is eventually the quantum effect in this picture: In the simple scale-up of QCD we would have

$$\text{Naturalness (QCD scale up)} : \quad m \sim \Lambda_\chi \sim \Lambda_{\text{TC}} \ll \Lambda_0. \quad (3)$$

The original version of TC, just a simple scale-up of QCD, however, is plagued by the notorious problems: Excessive flavor-changing neutral currents (FCNCs), and excessive oblique corrections of $\mathcal{O}(1)$ to the Peskin-Takeuchi S parameter³ compared with the typical experimental bound about 0.1.

The FCNC problem was resolved long time ago by the TC based on the near conformal gauge dynamics with $\gamma_m \simeq 1$,^{4,5} initially dubbed “scale-invariant TC” and then “walking TC”, with almost *non-running (conformal) gauge coupling*, based on the pioneering work by Maskawa and Nakajima¹ who discovered *non-zero critical coupling*, $\alpha_{\text{cr}} (\neq 0)$, for the $S\chi\text{SB}$ to occur. We may call it “Scale-invariant/Walking/Conformal TC” (SWC TC) (For reviews see Ref.⁶).

In addition to solving the FCNC problem, the theory made a definite prediction of “Techni-dilaton (TD)”,⁴ a pseudo Nambu-Goldstone (NG) boson of the spontaneous breaking of the (approximate) scale invariance of the theory. This will be the main topics of this talk in the light of modern version of SWC TC.

The modern version^{7–9} of SWC TC is based on the Caswell-Banks-Zaks (CBZ) infrared (IR) fixed point¹⁰, $\alpha_* = \alpha_*(N_f, N_{\text{TC}})$, which appears at two-loop beta function for the number of massless flavors $N_f (< 11N_{\text{TC}}/2)$ larger than a certain number $N_f^* (\gg N_{\text{TC}})$. See Fig. 1 and later discussions. Due to the IR fixed point the




Fig. 1. The beta function and $\alpha(\mu)$ for SWC-TC.

coupling is almost non-running (“walking”) all the way up to the intrinsic scale Λ_{TC} which is generated by the scale anomaly associated with the (two-loop) running of the coupling analogously to QCD scale-up in Eq. (2). For $\mu > \Lambda_{\text{TC}}$ (Region I of Fig. 1) the coupling no longer walks and runs similarly to that of QCD. When we set α_* slightly larger than α_{cr} , we have a condensate or the dynamical mass of the techni-fermion $m (\sim \Lambda_\chi)$, much smaller than the intrinsic scale of the theory $m \ll \Lambda_{\text{TC}}$. The CBZ-IR fixed point α_* actually disappears (then becoming would-be IR fixed point) at the scale $\mu \lesssim m$ where the techni-fermions have acquired the mass m and get decoupled from the beta function for $\mu < m$ (Region III in Fig. 1). Nevertheless, the coupling is still walking due to the remnant of the CBZ-IR fixed point conformality in a wide region $m < \mu < \Lambda_{\text{TC}}$ (Region II in Fig. 1). Thus the *symmetry responsible for the natural hierarchy $m \sim \Lambda_\chi \ll \Lambda_{\text{TC}}$ is the (approximate) conformal symmetry*, while the naturalness for the hierarchy $\Lambda_{\text{TC}} \ll \Lambda_0$ is the same as that of QCD scale-up in Eq.(3):

$$\text{Naturalness (SWC TC)} : \quad m \sim \Lambda_\chi \ll \Lambda_{\text{TC}} \quad (\ll \Lambda_0). \quad (4)$$

The theory acts like the SWC-TC^{4,5} : It develops a large anomalous dimension $\gamma_m \simeq 1$ for the almost non-running coupling in the Region II.^{8,9} Here Λ_{TC} plays a

role of cutoff Λ identified with the ETC scale: $\Lambda_{\text{TC}} = \Lambda = \Lambda_{\text{ETC}}$.

Moreover, there also exists a possibility^{11,12} that the S parameter can be reduced in the case of SWC-TC.

In this talk I will argue^{13 a} that in contrast to the simple QCD scale-up which is widely believed to have no composite Higgs particle (“higgsless”), a salient feature of SWC TC is the *conformality which manifests itself by the appearance of a composite Higgs boson (“conformal Higgs”) as the Techni-dilaton (TD)*⁴ with mass relatively lighter than other techni-hadrons: $M_{\text{TD}} < M_\rho, M_{a_1} \cdots = \mathcal{O}(\Lambda_\chi) \ll \Lambda_{\text{TC}} = \Lambda_{\text{ETC}}$, where $M_\rho, M_{a_1} \cdots$ denote the mass of techni- ρ , techni- a_1 , etc. This is contrasted to the QCD dynamics where there are no scalar bound states lighter than others. Note that there is no idealized limit where the TD becomes exactly massless to be a true NG boson, in sharp contrast to the chiral symmetry breaking. Scale symmetry is always broken explicitly as well as spontaneously^b.

For the phenomenological purpose, I will argue through several different calculations^{13,15,16} that the techni-dilaton mass in the typical SWC TC models will be in the range (see the footnote below Eq.(30), however):

$$m_{\text{TD}} = 500 - 600 \text{ GeV}, \quad (5)$$

which is definitely larger than the SM Higgs bound but still within the discovery region of the LHC experiments.

2. Scale-invariant/Walking/Conformal Technicolor

Let us briefly review the SWC TC.

The FCNC problem is related with the mass generation of quarks/leptons mass. In order to communicate the techni-fermion condensate to the quarks/leptons masses $m_{q/l}$, we would need interactions between the quarks/leptons and the techni-fermions which are typically introduced through Extended TC (ETC)^{17 c} with much higher scale $\Lambda_{\text{ETC}} (\gg \Lambda_\chi)$: $m_{q/l} \sim \frac{-1}{\Lambda_{\text{ETC}}^2} \langle \bar{T}T \rangle_{\Lambda_{\text{ETC}}}$, where $\langle \bar{T}T \rangle_{\Lambda_{\text{ETC}}}$ is the condensate measured at the scale of Λ_{ETC} . (We here do not refer to the origin of the mass scale Λ_{ETC} which should also be of dynamical origin such as the tumbling.) Since the newly introduced ETC interactions characterized by the same scale Λ_{ETC} should induce extra FCNC's, we should impose a constraint $\Lambda_{\text{ETC}} > 10^6 \text{ GeV}$ in order to avoid the excessive FCNC's (typically involving s quark). If we assume a

^aPreliminary discussions on the revival of the techni-dilaton⁴ were given in several talks¹⁴.

^bThe straightforward calculations near the conformal edge indicated¹⁶ that there is no isolated massless spectrum: $M_{\text{TD}}/F_\pi, M_{\text{TD}}/M_\rho, \cdots \rightarrow \text{const.} \neq 0$ even in the limit of $\alpha_* \rightarrow \alpha_{\text{cr}} (N_f \rightarrow N_f^{\text{crit}})$ where $F_\pi/\Lambda_{\text{TC}}, M_{\text{TD}}/\Lambda_{\text{TC}}, M_\rho/\Lambda_{\text{TC}}, \cdots \rightarrow 0$. In the case of holographic TD,¹³ this fact is realized in a different manner: Although there apparently exists an isolated massless spectrum, $M_{\text{TD}}/F_\pi \rightarrow 0$ while $M_\rho/F_\pi, M_{a_1}/F_\pi \rightarrow \text{const.} \neq 0$, the decay constant of the TD diverges $F_{\text{TD}}/F_\pi \rightarrow \infty$ in that limit and hence it gets decoupled. See later discussions.

^cThe same can be done in a composite model where quarks/leptons and techni-fermions are composites on the same footing.¹⁸

simple QCD scale up, $\langle \bar{T}T \rangle_{\Lambda_{\text{ETC}}} \simeq \langle \bar{T}T \rangle_{\Lambda_\chi} = -N_{\text{TC}} \cdot \Lambda_\chi^3$, we would have

$$m_{q/l} \sim \frac{\Lambda_\chi^3}{\Lambda_{\text{ETC}}^2} \cdot N_{\text{TC}} < 0.1 \text{ MeV} \cdot N_{\text{TC}} \left(\frac{N_c/N_{\text{TC}}}{N_d} \right)^{3/2}, \quad (6)$$

which implies that the typical mass (s -quark mass) would be roughly 10^{-3} smaller than the reality. We would desperately need 10^3 times enhancement.

This was actually realized dynamically by the TC based on the near conformal gauge dynamics,^{4,5} based on the Maskawa-Nakajima solution¹ of the (scale-invariant) ladder Schwinger-Dyson (SD) equation for fermion full propagator $S_F(p)$ parameterized as $iS_F^{-1}(p) = A(p^2)\not{p} - B(p^2)$ with *non-running* (conformal, an ideal limit of the “walking”) gauge coupling, $\alpha(Q) \equiv \alpha = \text{constant}$, with $Q^2 \equiv -p^2 > 0$. (See Fig. 2)

Maskawa and Nakajima discovered that the $S_\chi\text{SB}$ can only take place for strong coupling $\alpha > \alpha_{\text{cr}} = \mathcal{O}(1)$, *non-zero critical coupling*.^d The critical value reads:²⁰ $C_2(F)\alpha_{\text{cr}} = \pi/3$, or

$$\alpha_{\text{cr}} = (\pi/3) \cdot 2N_{\text{TC}}/(N_{\text{TC}}^2 - 1) \quad (7)$$

in the $SU(N_{\text{TC}})$ gauge theory, where $C_2(F)$ is the quadratic Casimir of the technifermion representation of the TC. The asymptotic form of the Maskawa-Nakajima $S_\chi\text{SB}$ solution of the fermion mass function $\Sigma(Q) = B(p^2)/A(p^2)$ in Landau gauge ($A(p^2) \equiv 1$) reads,^{1,20}

$$\Sigma(Q) \sim 1/Q \quad (Q \gg \Lambda_\chi). \quad (8)$$

We then proposed a “Scale-invariant TC”⁴, based on the observation that Eq.(8) implies a special value of the anomalous dimension

$$\gamma_m = -\Lambda \frac{\partial \ln Z_m}{\partial \Lambda} = 1, \quad (9)$$

to be compared with the operator product expansion (OPE), $\Sigma(Q) \sim 1/Q^2 \cdot (Q/\Lambda_\chi)^{\gamma_m}$. Accordingly, we had an enhanced condensate $\langle \bar{T}T \rangle_{\Lambda_{\text{ETC}}} = Z_m^{-1} \cdot \langle \bar{T}T \rangle_{\Lambda_\chi} \simeq -N_{\text{TC}}(\Lambda_{\text{ETC}}\Lambda_\chi^2)$, with the (inverse) mass renormalization constant being $Z_m^{-1} = (\Lambda_{\text{ETC}}/\Lambda_\chi)^{\gamma_m} \simeq \Lambda_{\text{ETC}}/\Lambda_\chi \simeq 10^3$, which in fact yields the desired enhancement. We actually obtained a different formula than Eq.(6).⁴

$$m_{q/l} \sim \frac{\Lambda_\chi^2}{\Lambda_{\text{ETC}}} \cdot N_{\text{TC}}. \quad (10)$$




Fig. 2. Graphical expression of the SD equation in the ladder approximation.

^d Earlier works¹⁹ in the ladder SD equation with non-running coupling all confused explicit breaking solution with the SSB solution and thus implied $\alpha_{\text{cr}} = 0$

The model⁴ was formulated in terms of the Renormalization Group Equation (RGE) a la Miransky²¹ for the Maskawa-Nakajima solution $\Sigma(m) = m$ which takes the form^{20,21}

$$\Lambda_\chi \sim m \sim 4\Lambda \exp\left(-\pi/\sqrt{\alpha/\alpha_{\text{cr}} - 1}\right), \quad (11)$$

where Λ is the cutoff of the SD equation. This has an *essential singularity* often called “Miransky scaling” and implies the *non-perturbative beta function* having a *multiple zero*^e:

$$\beta(\alpha)_{\text{NP}} = \Lambda \frac{\partial \alpha(\Lambda)}{\partial \Lambda} = -\frac{2}{3C_2(F)} \left(\frac{\alpha}{\alpha_{\text{cr}}} - 1\right)^{3/2}, \quad (12)$$

with the critical coupling α_{cr} identified with a nontrivial ultraviolet (UV) stable fixed point $\alpha = \alpha(\Lambda) \rightarrow \alpha_{\text{cr}}$ as $\Lambda/m \rightarrow \infty$.

Subsequently, similar enhancement effects of the condensate were also studied⁵ within the same framework of the ladder SD equation, without use of the RGE concepts of anomalous dimension and fixed point, rather emphasizing the asymptotic freedom of the TC theories with slowly-running (walking) coupling which was implemented into the ladder SD equation (“improved ladder SD equation”).

Today the Scale-invariant/Walking/Conformal TC (SWC TC) is simply characterized by near conformal property with $\gamma_m \simeq 1$ (For a review see Ref.⁶). Such a theory should have an almost non-running and strong gauge coupling (larger than a certain non-zero critical coupling for S χ SB) to be realized either at UV fixed point or IR fixed point, or both (“fusion” of the IR and UV fixed points), as was characterized by “*Conformal Phase Transition (CPT)*”.⁹

The essential feature of the above is precisely what happens in the modern version⁷⁻⁹ of the SWC TC based on the CBZ IR fixed point¹⁰ of the large N_f QCD, the QCD-like theory with many flavors $N_f (\gg N_{\text{TC}})$ of massless techni-fermions,^f see Fig. 1. The two-loop beta function is given by $\beta(\alpha) = \mu \frac{d}{d\mu} \alpha(\mu) = -b\alpha^2(\mu) - c\alpha^3(\mu)$, where $b = (11N_{\text{TC}} - 2N_f)/(6\pi)$, $c = [34N_{\text{TC}}^2 - 10N_f N_{\text{TC}} - 3N_f(N_{\text{TC}}^2 - 1)/N_{\text{TC}}]/(24\pi^2)$. When $b > 0$ and $c < 0$, i.e., $N_f^* < N_f < \frac{11}{2}N_{\text{TC}}$ ($N_f^* \simeq 8.05$ for $N_{\text{TC}} = 3$), there exists an IR fixed point (CBZ IR fixed point) at $\alpha = \alpha_*$, $\beta(\alpha_*) = 0$, where

$$\alpha_* = \alpha_*(N_{\text{TC}}, N_f) = -b/c. \quad (13)$$

Note that $\alpha_* = \alpha_*(N_f, N_{\text{TC}}) \rightarrow 0$ as $N_f \rightarrow 11N_{\text{TC}}/2$ ($b \rightarrow 0$) and hence there exists a certain range $N_f^{\text{cr}} < N_f < 11N_{\text{TC}}/2$ (“Conformal Window”) satisfying $\alpha_* < \alpha_{\text{cr}}$, where the gauge coupling $\alpha(\mu) (< \alpha_*)$ gets so weak that attractive forces are no longer strong enough to trigger the S χ SB as was demonstrated by Maskawa-

^e Simple zero of the beta function, $\beta(\alpha) \sim (\alpha - \alpha_{\text{cr}})^1$, never reproduces the essential singularity scaling, as is evident from Eq.(2).

^f For SWC TC based on higher representation/other gauge groups see, e.g., Ref.²²

Nakajima.¹ N_f^{cr} such that $\alpha_*(N_{\text{TC}}, N_f^{\text{cr}}) = \alpha_{\text{cr}}$ may be evaluated by using the value of α_{cr} from the ladder SD equation Eq.(7):⁸ $N_f^{\text{cr}} \simeq 4N_{\text{TC}}$ ($= 12$ for $N_{\text{TC}} = 3$)⁸.

Here we are interested in the $S\chi\text{SB}$ phase slightly off the conformal window, $0 < \alpha_* - \alpha_{\text{cr}} \ll 1$ ($N_f \simeq N_f^{\text{cr}}$). We may use the same equation as the ladder SD equation with $\alpha(\mu) \simeq \text{const.} = \alpha_*$, yielding the same form as Eq.(11):⁸

$$m \sim 4\Lambda_{\text{TC}} \exp\left(-\pi/\sqrt{\alpha_*/\alpha_{\text{cr}} - 1}\right) \ll \Lambda_{\text{TC}} \quad (\alpha_* \simeq \alpha_{\text{cr}}), \quad (14)$$

where the cutoff Λ was identified with $\Lambda_{\text{TC}} (= \Lambda_{\text{ETC}})$. We also have the same result as Eqs. (8),(9):

$$\Sigma(Q) \sim 1/Q, \quad \gamma_m \simeq 1. \quad (15)$$

Hence it acts like SWC TC. Incidentally, Eq.(14) implies a *multiple zero* at $\alpha_* = \alpha_{\text{cr}}$ in a non-perturbative beta function for $\alpha_* = \alpha_*(\Lambda)$ similar to Eq.(12), which would suggest “running” of the IR fixed point α_* with its UV fixed point $\alpha_* = \alpha_{\text{cr}}$ in the limit $\Lambda_{\text{TC}}/m \rightarrow \infty$.

The actual running of the coupling largely based on two-loop perturbation is already depicted in Fig. 1. The critical coupling α_{cr} can be regarded as the UV fixed point viewed from the IR part of the Region II ($m < \mu < \mu_{\text{cr}}$, with μ_{cr} such that $\alpha(\mu_{\text{cr}}) = \alpha_{\text{cr}}$), while it is regarded as the IR fixed point from the UV part of the Region II ($\Lambda_{\text{TC}} > \mu > \mu_{\text{cr}}$), with the Region II regarded as the *fusion of the IR and UV fixed points* in the idealized limit of non-running (perturbative) coupling in Region II (or $\Lambda_{\text{TC}}/m \rightarrow \infty$). Although the perturbative (two-loop) beta function has a *simple zero*, which *never corresponds to the essential singularity* scaling as we noted before, the coupling near α_{cr} should be sensitive to the non-perturbative effects in such a way that the beta function looks like the *multiple zero* as in Eq. (12) from both sides, corresponding to the *essential singularity scaling* as in Eq. (11). This should be tested by the fully non-perturbative studies like lattice simulations. A possible phase diagram (Fig 3 of Ref.⁹) of the large N_f QCD on the lattice is also waiting for the test by simulations.

3. Conformal Phase Transition^{9,14}

Such an *essential singularity* scaling law like Eq.(11),(14), or equivalently the *multiple zero* of the non-perturbative beta function, characterizes an unusual phase transition, what we called “*Conformal Phase Transition (CPT)*”, where the Ginzburg-Landau effective theory breaks down:⁹ Although it is a second order (continuous) phase transition where the order parameter m ($\alpha_* > \alpha_{\text{cr}}$) is continuously changed to $m = 0$ in the symmetric phase (conformal window, $\alpha_* < \alpha_{\text{cr}}$), the spectra do not, i.e., while there exist light composite particles whose mass vanishes at the critical

⁸ The value should not be taken seriously, since $\alpha_* = \alpha_{\text{cr}}$ is of $\mathcal{O}(1)$ and the perturbative estimate of α_* is not so reliable there, although the chiral symmetry restoration in large N_f QCD has been supported by many other arguments, most notably the lattice QCD simulations,^{23,24} which however suggest diverse results as to N_f^{cr} ; See e.g.,²⁵ for recent results.

point when approached from the side of the SSB phase, no isolated light particles do not exist in the conformal window, recently dubbed “unparticle”.²⁶ This reflects the feature of the conformal symmetry in the conformal window. In fact explicit computations show no light (composite) spectra in the conformal window, in sharp contrast to the S χ SB phase where light composite spectra do exist with mass of order $\mathcal{O}(m)$ which vanishes as we approach the conformal window $N_f \nearrow N_f^{\text{cr}}$.^{8,9,16}

The essence of CPT was illustrated⁹ by a simpler model, 2-dimensional Gross-Neveu Model. This is the $D \rightarrow 2$ limit of the D -dimensional Gross-Neveu model ($2 < D < 4$) which has the beta function and the anomalous dimension:^{27,28}

$$\beta(g) = -2g(g - g_*), \quad \gamma_m = 2g, \quad (16)$$

where $g = g_*(\equiv D/2 - 1) = g_{\text{cr}}$ and $g = 0$ are respectively the UV and IR fixed points of the dimensionless four-fermion coupling, g , properly normalized (as $g_* = 1$ for the $D = 4$ NJL model). There exist light composites π, σ near the UV fixed point (phase boundary) $g \simeq g_*$ in both sides of symmetric ($0 < g < g_*$) and SSB ($g > g_*$) phases as in the NJL model.

Now we consider $D \rightarrow 2$ ($g_* \rightarrow 0$) where we have a well-known effective potential: $V(\sigma, \pi) \sim (1/g - 1)\rho^2 + \rho^2 \ln(\rho^2/\Lambda^2)$, or $\partial^2 V/\partial \rho^2|_{\rho=0} = -\infty$, where $\rho^2 = \pi^2 + \sigma^2$. This implies breakdown of the Ginzburg-Landau theory which distinguishes the SSB (< 0) and symmetric (> 0) phases by the signature of the finite $\partial^2 V/\partial \rho^2$ at the critical point $g = 0$. Eq. (16) now reads:

$$\beta(g) = -2g^2, \quad \gamma_m|_{g=0} = 0 \quad (D = 2), \quad (17)$$

namely a *fusion of the UV and IR fixed points* at $g = 0$ as a result of *multiple zero* (not a simple zero) at $g = 0$. Now the symmetric phase is squeezed out to the region $g < 0$ (conformal phase) which corresponds to a *repulsive* four-fermion interaction and no composite states exist, while in the SSB phase ($g > 0$) there exists a composite state σ of mass $M_\sigma = 2m$ where the dynamical mass of the fermion is given by $m^2 \sim \Lambda^2 \exp(-1/g) \rightarrow 0$ ($g \rightarrow +0$), which shows an *essential singularity* scaling, in accord with the beta function with *multiple zero*, $\beta(g) = \Lambda \partial g/\partial \Lambda = -2g^2$. Note the would-be composite mass in the symmetric phase $|M|^2 \sim \Lambda^2 \exp(-1/g) \rightarrow \infty$ ($g \rightarrow -0$).

Now look at the SWC TC as modeled by the large N_f QCD: When the walking coupling $\alpha(Q) \simeq \alpha_*$ is close to the critical coupling, $\alpha_* \simeq \alpha_{\text{cr}}$, we should include the *induced* four-fermion interaction, $(G/2)[(\bar{\psi}\psi)^2 + (\bar{\psi}i\gamma_5\psi)^2]$, which becomes relevant operator due to the anomalous dimension $\gamma_m = 1$, and the system becomes “gauged Nambu-Jona-Lasinio” model²⁹ whose solution in the full parameter space was obtained in Ref.³⁰

Thus we may regard the *SWC TC as the gauged Nambu-Jona-Lasinio (NJL) model*. It was found³⁰ that S χ SB solution exists for the parameter space $g > g_{(+)} = (1 + \sqrt{1 - \alpha_*/\alpha_{\text{cr}}})^2/4$ ($\alpha_* < \alpha_{\text{cr}}$) as well as the region $\alpha_* > \alpha_{\text{cr}}$, where the dimensionless four-fermion coupling $g \equiv G\Lambda^2 (N_{\text{TC}}/4\pi^2)$ is normalized as $g = 1$ for $\alpha_* = 0$ (pure NJL model without gauge interaction). Based on the solution (including the

running coupling case), the RGE flow in (α, g) space was found to be along the line of $\alpha = \alpha_*$ (α does not run),^h on which the four-fermion coupling g runs, with the beta function and anomalous dimension given by^{28,32,33}

$$\beta(g) = -2(g - g_{(+)})(g - g_{(-)}), \quad \gamma_m = 2g + \alpha_*/(2\alpha_{\text{cr}}) \quad (18)$$

where $g = g_{(\pm)} \equiv (1 \pm \sqrt{1 - \alpha_*/\alpha_{\text{cr}}})^2/4$ are regarded as the UV/IR fixed points (fixed lines) for $\alpha_* \leq \alpha_{\text{cr}}$. The above anomalous dimension takes the values: $\gamma_m = 1 + \sqrt{1 - \alpha_*/\alpha_{\text{cr}}}$ ³¹ at the UV fixed line while $\gamma_m = 1 - \sqrt{1 - \alpha_*/\alpha_{\text{cr}}}$ at the IR fixed line. Light composite spectra only exist near the UV fixed line (phase boundary) $g \simeq g_{(+)}$ in both SSB ($g > g_{(+)}$) and symmetric ($g < g_{(+)}$) phases as in NJL model. Thus it follows that as $\alpha_* \rightarrow \alpha_{\text{cr}}$ Eq. (18) takes the form

$$\beta(g) = -2(g - g_*)^2, \quad \gamma_m|_{g=g_*} = 1, \quad (\alpha_* = \alpha_{\text{cr}}), \quad (19)$$

with $g_{(\pm)} \rightarrow 1/4 \equiv g_*$, and hence we again got a *multiple zero* and *fusion of UV and IR fixed lines*^{28,32,33} which corresponds to the *essential singularity* scaling;³⁰ $m^2 \sim \Lambda^2 \exp(-1/(g - g_*))$. A similar observation was also made recently.³⁴

In passing, it should be stressed that the *anomalous dimension never changes discontinuously across the phase boundary* as is seen from Eq.(16) and Eq.(18).^{27,28,32}

The scale anomaly in this case is given by:⁹

$$\begin{aligned} \langle \partial^\mu D_\mu \rangle &= \langle \theta_\mu^\mu \rangle = 4\langle \theta_0^0 \rangle = \frac{\beta(g)}{g} \cdot \frac{G}{2} \langle (\bar{\psi}\psi)^2 + (\bar{\psi}i\gamma_5\psi)^2 \rangle \\ &\simeq -m^4 \cdot (4N_f N_{\text{TC}}/\pi^4) = \mathcal{O}(\Lambda_\chi^4), \end{aligned} \quad (20)$$

where the second line was from the explicit computation³⁵ of the vacuum energy $\langle \theta_0^0 \rangle$ in the limit $\Lambda/m \rightarrow \infty$ ($g \rightarrow g_*$) at $\alpha \equiv \alpha_{\text{cr}}$ (The result coincides with the one for $\alpha \rightarrow \alpha_{\text{cr}}$ with $g \equiv 0$, see Eq.(24).³⁶). Again there is a composite state this time having mass¹⁵

$$M_\sigma \rightarrow \sqrt{2}m, \quad (21)$$

as $g \rightarrow g_* + 0$, while there are no composites $|M|^2 \sim \Lambda^2 \exp(-1/(g - g_*)) \rightarrow \infty$ for $g \rightarrow g_* - 0$. Eq.(21) is compared with $M_\sigma = 2m$ in the pure NJL case with $\alpha \equiv 0$. This slightly lighter scalar may be identified with the techni-dilaton in the SWC TC. I will come back to this later.

The absence of the composites in the symmetric phase $g < g_*$ may be understood as in the 2-dimensional Gross-Neveu model for $g < 0$, namely the *repulsive* four-fermion interactions: From the analysis of the RG flow, it was argued³² that the IR fixed line $g = g_{(-)}$ is due to the *induced* four-fermion interaction by the walking TC dynamics itself, while deviation from that line, $g - g_{(-)}$, is due to the *additional* four-fermion interactions, repulsive ($g < g_{(-)}$) and attractive ($g > g_{(-)}$), from UV

^h The beta function in Eq.(12) may be regarded as an artificial one keeping $g \equiv \text{const.}$ which is not along the renormalized trajectory in the extended parameter space (α, g) .

dynamics other than the TC (i.e., ETC). It is clear that no light composites exist for repulsive four-fermion interaction $g < g_{(-)}$, which becomes $g < g_*$ at $\alpha_* = \alpha_{\text{cr}}$.

4. S Parameter Constraint

Now we come to the next problem of TC, so-called S, T, U parameters³ measuring possible new physics in terms of the deviation of the LEP precision experiments from the SM. In particular, S parameter excludes the TC as a simple scale-up of QCD which yields $S = (N_f/2) \cdot \hat{S}$ with $\hat{S}_{\text{QCD}} = 0.32 \pm 0.04$. For a typical ETC model with one-family TC, $N_f = 8$,² we would get $S = \mathcal{O}(1)$ which is much larger than the experiments $S < 0.1$. This is the reason why many people believe that the TC is dead. However, since the simple scale-up of QCD was already ruled out by the FCNC as was discussed before, the real problem is whether or not the walking/conformal TC which solved the FCNC problem is also consistent with the S parameter constraint above. There have been many arguments^{11,12} that the S parameter value could be reduced in the walking/conformal TC than in the simple scale-up of QCD. Recently such a reduction has also been argued^{37,38} in a version of the holographic QCD³⁹ deformed to the walking/conformal TC by tuning a parameter to simulate the large anomalous dimension $\gamma_m \simeq 1$.

Here we present the most straightforward computation of the S parameter for the large N_f QCD, based on the SD equation and (inhomogeneous) Bethe-Salpeter (BS) equation in the ladder approximation.¹² The S parameter $S = (N_f/2)\hat{S}$ is defined by the slope of the the current correlators $\Pi_{JJ}(Q^2)$ at $Q^2 = 0$:

$$\hat{S} = -4\pi \frac{d}{dQ^2} [\Pi_{VV}(Q^2) - \Pi_{AA}(Q^2)] \Big|_{Q^2=0}, \quad (22)$$

where $\delta^{ab}(q_\mu q_\nu/q^2 - g_{\mu\nu}) \Pi_{JJ}(q^2) = \mathcal{F.T.} \ i \langle 0 | T J_\mu^a(x) J_\nu^b(0) | 0 \rangle$, $(J_\mu^a(x) = V_\mu^a(x), A_\mu^a(x))$, with $F_\pi^2 = \Pi_{VV}(0) - \Pi_{AA}(0)$. The current correlators are obtained by closing the fermion legs of the BS amplitudes $\chi_\mu^{(J)}(p; q) \sim \mathcal{F.T.} \ \langle 0 | T \psi(r/2) \bar{\psi}(-r/2) J_\mu(x) | 0 \rangle$, which is determined by the ladder BS equation (Fig.3). Solving the BS equation with the fermion propagator given as the solution




Fig. 3. Graphical expression of the BS equation in the ladder approximation.

of the ladder SD equation, we can evaluate the $\Pi_{VV}(Q^2) - \Pi_{AA}(Q^2)$ numerically. From this result we may read its slope at $Q^2 = 0$ to get $\hat{S}$.

The results show definitely smaller values of $\hat{S}$ than that in the ordinary QCD and moreover there is a tendency of the $\hat{S}$ getting reduced when approaching the

conformal window $\alpha_* \searrow \alpha_{\text{cr}} (N_f \nearrow N_f^{\text{cr}})$. However, due to technical limitation of the present computation getting very close to the conformal window, the reduction does not seem to be so dramatic as the walking TC being enough to be consistent with the experimental constraints. It is highly desirable to extend the computation further close to the conformal window.

Another approach to this problem is the deformation of the holographic QCD by the anomalous dimension. The reduction of S parameter in the SWC TC has been argued in a version of the hard-wall type bottom up holographic QCD³⁹ deformed to the SWCTC by tuning a parameter to simulate the large anomalous dimension $\gamma_m \simeq 1$.³⁷ We examined³⁸ such a possibility paying attention to the renormalization point dependence of the condensate. We explicitly calculated the S parameter in entire parameter space of the holographic SWC TC. We here take a set of F_π/M_ρ and γ_m . We find that $S > 0$ and it monotonically decreases to zero in accord with the previous results³⁷. However, our result turned out fairly independent of the value of the anomalous dimension γ_m , yielding no particular suppression solely by tuning the anomalous dimension large, $\hat{S} \sim B(F_\pi/M_\rho)^2 \rightarrow 0$ as $F_\pi/M_\rho \rightarrow 0$, with $B \simeq 27(32)$ for $\gamma_m \simeq 1(0)$, in sharp contrast to the previous claim³⁷. Although B contains full contributions from the infinite tower of the vector/axial-vector Kaluza-Klein modes (gauge bosons of hidden local symmetries)⁴⁰ of the 5-dimensional gauge bosons, the resultant value of B turned out close to $B \simeq 4\pi a \simeq 8\pi$ of the single ρ meson dominance, where $a \simeq 2$ is the parameter of the hidden local symmetry only for the ρ meson.⁴⁰ This implies that as far as the pure TC dynamics (without ETC dynamics, etc.) is concerned, an obvious way to dynamically reduce S parameter is to tune F_π/M_ρ very small, namely techni- ρ mass very large to several TeV region. (See, however, footnote below Eq.(30).)

We would need more dynamical information other than the holographic recipe, since the parameter corresponding to F_π/M_ρ as well as the scale parameter is a pure input in all the holographic models, whether bottom up or top down approach, in contrast to the underlying gauge theory which has only a single parameter, a scale parameter like Λ_{QCD} .

Curiously enough, when we calculate F_π/M_ρ from the SD and the homogeneous BS equations¹⁶ and S from the SD and the inhomogeneous BS equation¹² both in the straightforward calculation in the ladder approximation, a set of the calculated values of $(F_\pi/M_\rho, S)$ lies on the line of the holographic result.³⁸

5. Techni-dilaton

Now we come to the discussions of Techni-dilaton (TD). Existence of two largely separated scales, $\Lambda_\chi \sim m$ and Λ_{TC} such that $\Lambda_\chi \ll \Lambda_{\text{TC}}$, is the most important feature of SWC-TC, in sharp contrast to the ordinary QCD with small number of flavors (in the chiral limit) where all the mass parameters like dynamical mass of quarks are of order of the single scale parameter of the theory Λ_{QCD} , $m \sim \Lambda_\chi \sim \Lambda_{\text{QCD}}$. See Fig. 1. The intrinsic scale Λ_{TC} is related with the scale anomaly

corresponding to the *perturbative* running effects of the coupling, with the ordinary two-loop beta function $\beta(\alpha)$ in the Region I, in the same sense as in QCD.

$$\langle \partial^\mu D_\mu \rangle = \langle \theta_\mu^\mu \rangle = \frac{\beta(\alpha)}{4\alpha^2} \langle \alpha G_{\mu\nu}^2 \rangle = \mathcal{O}(\Lambda_{\text{TC}}^4), \quad (23)$$

which implies that all the techni-gluon balls have mass of $\mathcal{O}(\Lambda_{\text{TC}})$.

On the other hand, the scale Λ_χ is related with totally different scale anomaly due to the dynamical generation of m ($\sim \Lambda_\chi$) which does exist even in the idealized case with non-running coupling $\alpha(\mu) \equiv \alpha(> \alpha_{\text{cr}})$ such as the Maskawa-Nakajima solution,¹ as was discussed some time ago.³⁶ Such an idealized case well simulates the dynamics of Region II of Fig. 1,^{8,9} with anomalous dimension $\gamma_m \simeq 1$ and $m \ll \Lambda_{\text{TC}}$ in the numerical calculations,¹⁶ with the *perturbative* coupling constant in Region II being almost constant slightly larger than α_{cr} , $\alpha_{\text{cr}} < \alpha(\mu) (< \alpha_*)$, for a wide infrared region. The coupling $\alpha \equiv \alpha_*$ in the “idealized Region II” actually runs *non-perturbatively* according to the *essential-singularity* scaling (Miransky scaling²¹) of mass generation, Eq.(11), with the *non-perturbative* beta function $\beta_{\text{NP}}(\alpha)$, Eq.(12), having a *multiple zero* at $\alpha = \alpha_{\text{cr}}$. Then the *non-perturbative* scale anomaly reads⁹

$$\langle \partial^\mu D_\mu \rangle_{\text{NP}} = \langle \theta_\mu^\mu \rangle_{\text{NP}} = \frac{\beta_{\text{NP}}(\alpha)}{4\alpha^2} \langle \alpha G_{\mu\nu}^2 \rangle_{\text{NP}} = -m^4 \cdot \frac{4N_f N_{\text{TC}}}{\pi^4} = -\mathcal{O}(\Lambda_\chi^4), \quad (24)$$

where $\langle \cdots \rangle_{\text{NP}}$ is the quantity with the perturbative contributions subtracted:³⁶ $\langle \cdots \rangle_{\text{NP}} \equiv \langle \cdots \rangle - \langle \cdots \rangle_{\text{Perturbative}}$. Eq.(24) coincides with Eq.(20) and $\langle \partial^\mu D_\mu \rangle_{\text{NP}} / \Lambda_{\text{TC}}^4$ vanishes with $\langle \partial^\mu D_\mu \rangle_{\text{NP}} / m^4 \rightarrow \text{const.} \neq 0$, when we approach the conformal window from the broken phase $\alpha_* \searrow \alpha_{\text{cr}}$ ($m/\Lambda_{\text{TC}} \rightarrow 0$). All the techni-fermion bound states have mass of order of m ,⁴¹ while there are no light bound states in the symmetric phase (conformal window) $\alpha_* < \alpha_{\text{cr}}$, a characteristic feature of the conformal phase transition.⁹ The TD is associated with the latter scale anomaly and should have mass on order of m ($\ll \Lambda_{\text{TC}}$).

5.1. Calculation from Gauged NJL model in the ladder SD equation¹⁵

More concretely, the mass of TD or scalar bound state in the SWC-TC was estimated in various methods: The first method¹⁵ was based on the the ladder SD equation for the gauged NJL model which well simulates^{8,9} the conformal phase transition in the large N_f QCD. The result was already given by Eq.(21):

$$M_{\text{TD}} \simeq \sqrt{2}m. \quad (25)$$

ⁱIn terms of the gauged NJL model mentioned in Section 3 this is the expression of the scale anomaly for $\alpha \rightarrow \alpha_{\text{cr}}$ with $g = \text{const.} = g_*$, in contrast to Eq.(20) for $g \rightarrow g_*$ with $\alpha = \text{const.} = \alpha_* = \alpha_{\text{cr}}$. Both yield the same vacuum energy and hence the same scale anomaly.

5.2. Straightforward Calculation from Ladder SD and BS equations

Also a straightforward calculation¹⁶ of mass of TD, the scalar bound state was made in the vicinity of the CBZ-IR fixed point in the large N_f QCD, based on the coupled use of the ladder SD equation and (*homogeneous*) BS equation lacking the first term in Fig. 3: All the bound states masses are $M = \mathcal{O}(m)$ and $M/\Lambda_{\text{TC}} \rightarrow 0$, when approaching the conformal window $\alpha_* \rightarrow \alpha_{\text{cr}} (N_f \rightarrow N_f^{\text{cr}})$ such that $m/\Lambda_{\text{TC}} \rightarrow 0$. Near the conformal window ($N_f \nearrow N_f^{\text{cr}}$) the calculated values are $M_\rho/F_\pi \simeq 11$, $M_{a_1}/F_\pi \simeq 12$ (near degenerate!). On the other hand, the scalar mass sharply drops near the the conformal window, $M_{\text{TD}}/F_\pi \searrow 4$, or

$$M_{\text{TD}} \searrow 1.5m \simeq \sqrt{2}m (< M_\rho, M_{a_1}). \quad (26)$$

Note that in this calculation $M_{\text{TD}}/F_\pi \rightarrow \text{const.} \neq 0$ and hence there is no isolated massless scalar bound states even in the limit $N_f \rightarrow N_f^{\text{cr}}$. The result is consistent with Eq.(25) and is contrasted to the ordinary QCD where the scalar mass is larger than those of the vector mesons (“higgsless”) within the same framework of ladder SD/BS equation approach. The result would imply

$$m_{\text{TD}} \simeq 500 \text{ GeV} \quad (27)$$

in the case of the one-family TC model with $F_\pi \simeq 125 \text{ GeV}$.

5.3. Holographic Techni-dilaton¹³

Recently, we have calculated¹³ mass of TD in an extension of the previous paper³⁸ on the hard-wall-type bottom-up holographic SWC-TC by including effects of (techni-) gluon condensation parameterized as

$$\Gamma \equiv \left(\frac{(\frac{1}{\pi} \langle \alpha G_{\mu\nu}^2 \rangle / F_\pi^4)}{(\frac{1}{\pi} \langle \alpha G_{\mu\nu}^2 \rangle / f_\pi^4)_{\text{QCD}}} \right)^{1/4} \quad (28)$$

through the bulk flavor/chiral-singlet scalar field Φ_X , in addition to the conventional bulk scalar field Φ dual to the chiral condensate.

The five-dimensional action is given by

$$S_5 = \int d^4x \int_\epsilon^{z_m} dz \sqrt{-g} \frac{1}{g_5^2} e^{c g_5^2 \Phi_X(z)} \left(-\frac{1}{4} \text{Tr} [L_{MN} L^{MN} + R_{MN} R^{MN}] \right. \\ \left. + \text{Tr} [D_M \Phi^\dagger D^M \Phi - m_\Phi^2 \Phi^\dagger \Phi] + \frac{1}{2} \partial_M \Phi_X \partial^M \Phi_X \right) \quad (29)$$

where the anti-de-Sitter space (AdS_5) with the curvature radius L of AdS_5 is described by the metric $ds^2 = g_{MN} dx^M dx^N = (L/z)^2 (\eta_{\mu\nu} dx^\mu dx^\nu - dz^2)$ with $\eta_{\mu\nu} = \text{diag}[1, -1, -1, -1]$, $g = \det[g_{MN}] = -(L/z)^{10}$; g_5 denotes the gauge coupling in five-dimension and c is the dimensionless coupling constant, and $L_M(R_M) = L_M^a(R_M^a) T^a$ with the generators of $SU(N_f)$ are normalized by $\text{Tr}[T^a T^b] = \delta^{ab}$; $L(R)_{MN} = \partial_M L(R)_N - \partial_N L(R)_M - i[L(R)_M, L(R)_N]$. The covariant derivative acting on Φ is defined as $D_M \Phi = \partial_M \Phi + i L_M \Phi - i \Phi R_M$.

The TD, a flavor-singlet scalar bound state of techni-fermion and anti-techni-fermion, will be identified with the lowest KK mode coming from the bulk scalar field Φ , not Φ_X . Thanks to the additional explicit bulk scalar field Φ_X , we naturally improve the matching with the OPE of the underlying theory (QCD and SWC-TC) for current correlators so as to reproduce gluonic $1/Q^4$ term, which is clearly distinguished from the same $1/Q^4$ terms from chiral condensate in the case of SWC-TC with $\gamma_m \simeq 1$. Our model with $\gamma_m = 0$ and $N_f = 3$ well reproduces the real-life QCD.

It is rather straightforward^{37,39} to compute masses of the techni- ρ meson (M_ρ), the techni- a_1 meson (M_{a_1}), while for that of the TD, flavor-singlet scalar meson (M_{TD}), we would need additional IR potential with quartic coupling λ to stabilize the $S\chi\text{SB}$ vacuum.⁴² Such an IR potential might be regarded as generated by techni-fermion loop effects and we naturally expect $\lambda \sim N_{\text{TC}}/(4\pi)^2$. The S parameter was also calculated through the current correlators by the standard way. We found general tendency of the dependence of the meson masses relative to F_π , (M_ρ/F_π , M_{a_1}/F_π , M_{TD}/F_π) on γ_m , S , and Γ .

We find a characteristic feature of the techni-dilaton mass related to the conformality of SWC-TC: For fixed S and γ_m , absolute values of (M_ρ/F_π) and (M_{a_1}/F_π) are not sensitive to Γ , although they get *degenerate for large* Γ . On the contrary, (M_{TD}/F_π) *substantially decreases as* Γ *increases*. Actually, in the formal limit $\Gamma \rightarrow \infty$ we would have (M_{TD}/F_π) $\rightarrow 0$ (This is contrast to the straightforward computation through ladder SD and BS equations mentioned before¹⁶). For fixed S and Γ , again (M_ρ/F_π) and (M_{a_1}/F_π) are not sensitive to γ_m , while (M_{TD}/F_π) substantially decreases as γ_m increases.

Particularly for the case of $\gamma_m = 1$, we study the dependence of the S parameter on (M_ρ/F_π) for typical values of Γ . It is shown that the techni-gluon contribution reduces the value of S about 10% in the region of $\hat{S} \lesssim 0.1$, although the general tendency is similar to the previous paper³⁸ without techni-gluon condensation: $\hat{S}$ decreases to zero monotonically with respect to (F_π/M_ρ). This implies (M_ρ/F_π) necessarily increases when $\hat{S}$ is required to be smaller.

To be more concrete, we consider a couple of typical models of SWC-TC with $\gamma_m \simeq 1$ and $N_{\text{TC}} = 2, 3, 4$ based on the CBZ-IRFP in the large N_f QCD. Using the non-perturbative conformal anomaly Eq.(24) together with the non-perturbative beta function Eq.(12) and Eq.(11), we find a concrete relation between Γ and ($\Lambda_{\text{ETC}}/F_\pi$): In the case of $N_{\text{TC}} = 3$ ($N_f = 4N_{\text{TC}}$) and $S \simeq 0.1$, we have $\Gamma \simeq 7$ for ($\Lambda_{\text{ETC}}/F_\pi$) = 10^4 – 10^5 (required by the FCNC constraint). Thanks to the large anomalous dimension $\gamma_m \simeq 1$ and large techni-gluon condensation $\Gamma \simeq 7$, we obtain a relatively light techni-dilaton with mass

$$M_{\text{TD}} \simeq 600 \text{ GeV} \quad (30)$$

compared with $M_\rho \simeq M_{a_1} \simeq 3.8 \text{ TeV}$ (almost degenerate). Eq. (30) is consistent with the perturbative unitarity of $W_L W_L$ scattering even for large M_ρ, M_{a_1} . Note that largeness of M_ρ and M_{a_1} is essentially determined by the requirement of $S = 0.1$

fairly independently of technigluon condensation.^j

The essential reason for the large Γ is due to the existence of the wide conformal region $F_\pi(\sim m) < \mu < \Lambda_{\text{ETC}}$ with $\Lambda_{\text{ETC}}/F_\pi = 10^4\text{--}10^5$, which yields the smallness of the beta function (see Eq.(12) and Eq. (11)) and hence amplifies the technigluon condensation compared with the ordinary QCD with $\Gamma = 1$. Actually, in the idealized (phenomenologically uninteresting) limit $\Lambda_{\text{ETC}}/F_\pi \rightarrow \infty$, we would have $\Gamma \rightarrow \infty$, which in turn would imply $M_{\text{TD}}/F_\pi \rightarrow 0$ as mentioned above.^k

To conclude, various methods predicted the mass of the techni-dilaton (“conformal Higgs”) in the range of $M_{\text{TD}} \simeq 500 - 600$ GeV, which is within reach of LHC discovery.

Since the SWC TC models are strong coupling theories and the ladder approximation/holographic calculations would be no more than a qualitative hint, more reliable calculations are certainly needed, including the lattice simulations, before drawing a definite conclusion about the physics predictions. Besides the phase diagram including the TC-induced/ETC-driven four-fermion couplings on the lattice, more reliable calculations such as the spectra as well as anomalous dimensions, non-perturbative beta functions, S parameter, etc. are highly desired.

Acknowledgments

We thank K. Haba, M. Harada, M. Hashimoto, M. Kurachi, and S. Matsuzaki for collaborations on the relevant topics and useful discussions. This work was supported by the JSPS Grant-in-Aid for Scientific Research(S) # 22224003 and by the Nagoya University Foundation.

References

1. T. Maskawa and H. Nakajima, Prog. Theor. Phys. **52** (1974), 1326; **54** (1975), 860.
2. S. Weinberg, Phys. Rev. D **13** (1976), 974; D **19** (1979), 1277; L. Susskind, Phys. Rev. D **20** (1979), 2619; See for a review of earlier literature, E. Farhi and L. Susskind, Phys. Rep. **74** (1981), 277.
3. M. E. Peskin and T. Takeuchi, Phys. Rev. Lett. **65** (1990), 964; Phys. Rev. D **46** (1992), 381; B. Holdom and J. Terning, Phys. Lett. B **247** (1990), 88; M. Golden and L. Randall, Nucl. Phys. B **361** (1991), 3.

^jThe calculated S parameter here was from the TC dynamics alone and could be drastically changed by incorporating contributions from the generation mechanism of the mass of SM fermions such as the strong ETC dynamics. For instance, the fermion delocalization⁴³ in the Higgsless models as a possible analogue of certain ETC effects in fact can cancel large positive S arising from the 5-dimensional gauge sector which corresponds to the pure TC dynamics. If it is the case in the explicit ETC model, then the overall mass scale of techni-hadrons including TD would be much lower than the above estimate down to, say, 300 GeV.

^k This does not mean existence of true (exactly massless) NG boson of the broken scale symmetry, since such a would-be NG boson gets decoupled in our case: The decay constant of techni-dilaton F_{TD} would diverge, $F_{\text{TD}}/F_\pi \rightarrow \infty$, in that limit through the PCDC relation $F_{\text{TD}}^2 = -4\langle\theta_\mu^\mu\rangle/M_{\text{TD}}^2 \sim m^4/M_{\text{TD}}^2 \sim F_\pi^4/M_{\text{TD}}^2$. The scale symmetry is broken explicitly as well as spontaneously.

4. K. Yamawaki, M. Bando and K. Matumoto, Phys. Rev. Lett. **56** (1986), 1335; M. Bando, K. Matumoto and K. Yamawaki, Phys. Lett. B **178** (1986), 308. See also M. Bando, T. Morozumi, H. So and K. Yamawaki, Phys. Rev. Lett. **59** (1987), 389. The solution of the FCNC problem by the large anomalous dimension was first considered by B. Holdom, based on a pure assumption of the existence of UV fixed point without explicit dynamics and hence without definite prediction of the value of the anomalous dimension. See B. Holdom, Phys. Rev. D **24** (1981), 1441.
5. T. Akiba and T. Yanagida, Phys. Lett. B **169** (1986), 432; T. W. Appelquist, D. Karabali and L. C. R. Wijewardhana, Phys. Rev. Lett. **57** (1986), 957; T. Appelquist and L. C. R. Wijewardhana, Phys. Rev. D **36** (1987), 568. For an earlier work on this line based on pure numerical analysis see B. Holdom, Phys. Lett. B **150** (1985), 301.
6. C. T. Hill and E. H. Simmons, Phys. Rept. **381** (2003), 235 [Erratum-ibid. **390** (2004), 553]; K. Yamawaki, Lecture at 14th Symposium on Theoretical Physics, Cheju, Korea, July 1995, arXiv:hep-ph/9603293.
7. K. D. Lane and M. V. Ramana, Phys. Rev. D **44**, 2678 (1991).
8. T. Appelquist, J. Terning and L. C. Wijewardhana, Phys. Rev. Lett. **77** (1996), 1214; T. Appelquist, A. Ratnaweera, J. Terning and L. C. Wijewardhana, Phys. Rev. D **58** (1998), 105017.
9. V. A. Miransky and K. Yamawaki, Phys. Rev. D **55** (1997), 5051 [Errata; **56** (1997), 3768].
10. W. E. Caswell, Phys. Rev. Lett. **33**, 244 (1974); T. Banks and A. Zaks, Nucl. Phys. B **196** (1982), 189.
11. R. Sundrum and S. D. H. Hsu, Nucl. Phys. B **391** (1993), 127; T. Appelquist and G. Triantaphyllou, Phys. Lett. B **278** (1992), 345; T. Appelquist and F. Sannino, Phys. Rev. D **59** (1999), 067702.
12. M. Harada, M. Kurachi and K. Yamawaki, Prog. Theor. Phys. **115** (2006), 765 ; M. Kurachi and R. Shrock, Phys. Rev. D **74**, 056003 (2006); M. Kurachi, R. Shrock and K. Yamawaki, ibid D **76**, 035003 (2007).
13. K. Haba, S. Matsuzaki and K. Yamawaki, arXiv:1003.2841 [hep-ph]; arXiv:1006.2526 [hep-ph].
14. K. Yamawaki, Prog. Theor. Phys. Suppl. **180**, 1 (2010) [arXiv:0907.5277 [hep-ph]]; See also Prog. Theor. Phys. Suppl. **167**, 127 (2007).
15. S. Shuto, M. Tanabashi and K. Yamawaki, in *Proc. 1989 Workshop on Dynamical Symmetry Breaking*, Dec. 21-23, 1989, Nagoya, eds. T. Muta and K. Yamawaki (Nagoya Univ., Nagoya, 1990) 115-123; M. S. Carena and C. E. M. Wagner, Phys. Lett. B **285**, 277 (1992); M. Hashimoto, Phys. Lett. B **441**, 389 (1998).
16. M. Harada, M. Kurachi and K. Yamawaki, Phys. Rev. D **68** (2003), 076001; M. Kurachi and R. Shrock, JHEP **0612**, 034 (2006).
17. S. Dimopoulos and L. Susskind, Nucl. Phys. B **155** (1979), 237; E. Eichten and K. D. Lane, Phys. Lett. B **90** (1980), 125.
18. K. Yamawaki and T. Yokota, Phys. Lett. B **113** (1982), 293; Nucl. Phys. B **223** (1983), 144.
19. K. Johnson, M. Baker and R. Willey, Phys. Rev. **136** (1964), B1111; M. Baker and K. Johnson, Phys. Rev. **D3** (1971), 2516; R. Jackiw and K. Johnson Phys. Rev. **D8** (1973), 2386; J.M. Cornwall and R.E. Norton, ibid, 3338.
20. R. Fukuda and T. Kugo, Nucl. Phys. B **117** (1976), 250.
21. V. A. Miransky, Nuovo Cim. A **90** (1985), 149.
22. F. Sannino and K. Tuominen, Phys. Rev. D **71**, 051901 (2005); D. K. Hong, S. D. H. Hsu and F. Sannino, Phys. Lett. B **597**, 89 (2004); For a review F. Sannino, arXiv:0804.0182 [hep-ph].

23. Y. Iwasaki, K. Kanaya, S. Kaya, S. Sakai and T. Yoshie, Nucl. Phys. (Proc. Suppl.) **53** (1997), 449; Prog. Theor. Phys. Suppl. No. 131 (1998), 415.
24. T. Appelquist, G. T. Fleming and E. T. Neil, Phys.Rev.Lett.100:171607,2008, Erratum-ibid.102:149902,2009; Phys. Rev. D **79**, 076010 (2009); A. Deuzeman, M. P. Lombardo and E. Pallante, Phys. Lett. B **670**, 41 (2008); Talks presented at SCGT 09, <http://www.eken.phys.nagoya-u.ac.jp/scgt09/> .
25. Talks at SCGT 09: <http://www.eken.phys.nagoya-u.ac.jp/scgt09/> .
26. H. Georgi, Phys. Rev. Lett. **98**, 221601 (2007).
27. Y. Kikukawa and K. Yamawaki, Phys. Lett. B **234**, 497 (1990).
28. K. i. Kondo, M. Tanabashi and K. Yamawaki, Prog. Theor. Phys. **89**, 1249 (1993).
29. W. A. Bardeen, C. N. Leung and S. T. Love, Phys. Rev. Lett. **56**, 1230 (1986).
30. K. i. Kondo, H. Mino and K. Yamawaki, Phys. Rev. D **39**, 2430 (1989); K. Yamawaki, in *Proc. Johns Hopkins Workshop on Current Problems in Particle Theory 12, Baltimore, June 8-10, 1988*, edited by G. Domokos and S. Kovesi-Domokos (World Scientific Pub. Co., Singapore 1988); T. Appelquist, M. Soldate, T. Takeuchi, and L. C. R. Wijewardhana, *ibid.*
31. V. A. Miransky and K. Yamawaki, Mod. Phys. Lett. A **4** (1989), 129.
32. K. i. Kondo, S. Shuto and K. Yamawaki, Mod. Phys. Lett. A **6**, 3385 (1991).
33. K. I. Aoki, K. Morikawa, J. I. Sumi, H. Terao and M. Tomoyose, Prog. Theor. Phys. **102**, 1151 (1999).
34. D. B. Kaplan, J. W. Lee, D. T. Son and M. A. Stephanov, Phys. Rev. D **80**, 125005 (2009).
35. T. Nonoyama, T. B. Suzuki and K. Yamawaki, Prog. Theor. Phys. **81**, 1238 (1989).
36. V. A. Miransky and V. P. Gusynin, Prog. Theor. Phys. **81** (1989) 426.
37. D. K. Hong and H. U. Yee, Phys. Rev. D **74** (2006), 015011; M. Piai, arXiv:hep-ph/0608241; K. Agashe, C. Csaki, C. Grojean and M. Reece, JHEP **0712**, 003 (2007).
38. K. Haba, S. Matsuzaki and K. Yamawaki, Prog. Theor. Phys. **120**, 691 (2008).
39. J. Erlich, E. Katz, D. T. Son and M. A. Stephanov, Phys. Rev. Lett. **95** (2005), 261602; L. Da Rold and A. Pomarol, Nucl. Phys. B **721**, 79 (2005).
40. M. Bando, T. Kugo, S. Uehara, K. Yamawaki and T. Yanagida, Phys. Rev. Lett. **54** (1985), 1215; M. Bando, T. Kugo and K. Yamawaki, Nucl. Phys. B **259** (1985), 493; Phys. Rept. **164** (1988), 217; M. Harada and K. Yamawaki, Phys. Rept. **381** (2003), 1.
41. R. S. Chivukula, Phys. Rev. D **55** (1997), 5238.
42. L. Da Rold and A. Pomarol, JHEP **0601**, 157 (2006).
43. G. Cacciapaglia, C. Csaki, C. Grojean and J. Terning, Phys. Rev. D **71**, 035015 (2005); R. Foadi, S. Gopalakrishna and C. Schmidt, Phys. Lett. B **606**, 157 (2005). R. S. Chivukula, E. H. Simmons, H. J. He, M. Kurachi and M. Tanabashi, Phys. Rev. D **72**, 015008 (2005)